Chad J. Muxlow, D.O.

Director of Sports Medicine Surgery,
Sentara RMH Orthopedics and Sports Medicine
Head Team Surgeon, Bridgewater College
Head Team Surgeon, Eastern Mennonite University
www.MuxlowSportsMedicine.com



Physical Therapy Prescription – Arthroscopic Meniscectomy, Chondral Debridement

Name:	Date:
Procedure: R / L	Date of Surgery:
Frequency: 2-3 times per week for weeks	

PHASE I (Weeks 0 – 2): decrease edema, activate quadriceps

- Weightbearing: As tolerated; okay to use crutches for 2-3 days if needed
- Brace: None
- Range of Motion: AAROM → AROM as tolerated
- Therapeutic Exercises: Patellar mobs, quad/hamstring sets, heel slides, step-ups, straight-leg raises, stationary bike as tolerated; core exercises
- Modalities: Per therapist, including electrical stimulation, ultrasound, heat (before), ice (after)

Phase II (Weeks 2-4)

- Weightbearing: As tolerated
- Brace: None
- Range of Motion: Full
- Therapeutic Exercises: Progress Phase I exercises; lunges, wall-sits; add cycling and elliptical
- Modalities: Per therapist, including electrical stimulation, ultrasound, heat (before), ice (after)

Phase III (Weeks 4-6)

- Weightbearing: As tolerated
- Brace: None
- Range of Motion: Full
- Therapeutic Exercises: Progress Phase II exercises; add plyometrics and sport-specific exercises; add running; return to athletic activity as tolerated at week 6
- Modalities: Per therapist, including electrical stimulation, ultrasound, heat (before), ice (after)

Signature:	Date: